




Uniwersytet Rolniczy w Krakowie
University of Agriculture in Krakow
<https://repo.ur.krakow.pl>

Publikacja / Publication	Effect of water treatment with low-temperature and low-pressure glow plasma of low frequency on the growth of selected microorganisms, Chwastowski Jarosław, Wójcik Katarzyna, Kołoczek Henryk, Oszczęda Zdzisław , Khachatryan Karen, Tomasik Piotr
DOI wersji wydawcy / Published version DOI	http://dx.doi.org/10.1080/10942912.2023.2169708
Adres publikacji w Repozytorium URL / Publication address in Repository	https://repo.ur.krakow.pl/info/article/URK5c572bb81e4d4fca92843b7e49ec2804/
Data opublikowania w Repozytorium / Deposited in Repository on	30 sty 2023
Cytuj tę wersję / Cite this version	Chwastowski Jarosław, Wójcik Katarzyna, Kołoczek Henryk, Oszczęda Zdzisław , Khachatryan Karen, Tomasik Piotr: Effect of water treatment with low-temperature and low-pressure glow plasma of low frequency on the growth of selected microorganisms, International Journal of Food Properties, vol. 26, nr 1, 2023, s. 502-510, DOI:10.1080/10942912.2023.2169708



Effect of water treatment with low-temperature and low-pressure glow plasma of low frequency on the growth of selected microorganisms

Jarosław Chwastowski, Katarzyna Wójcik, Henryk Kołoczek, Zdzisław Oszczęda, Karen Khachatryan & Piotr Tomasiak

To cite this article: Jarosław Chwastowski, Katarzyna Wójcik, Henryk Kołoczek, Zdzisław Oszczęda, Karen Khachatryan & Piotr Tomasiak (2023) Effect of water treatment with low-temperature and low-pressure glow plasma of low frequency on the growth of selected microorganisms, *International Journal of Food Properties*, 26:1, 502-510, DOI: [10.1080/10942912.2023.2169708](https://doi.org/10.1080/10942912.2023.2169708)

To link to this article: <https://doi.org/10.1080/10942912.2023.2169708>



Published with license by Taylor & Francis Group, LLC. © 2023 Jarosław Chwastowski, Katarzyna Wójcik, Henryk Kołoczek, Zdzisław Oszczęda, Karen Khachatryan and Piotr Tomasiak



Published online: 27 Jan 2023.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Effect of water treatment with low-temperature and low-pressure glow plasma of low frequency on the growth of selected microorganisms

Jarosław Chwastowski^a, Katarzyna Wójcik^b, Henryk Kołoczek^a, Zdzisław Oszczyda^c, Karen Khachatryan^d, and Piotr Tomasik^c

^aInstitute of Chemistry and Inorganic Technology, Krakow University of Technology, Krakow, Poland; ^bCentral Laboratory for Diagnostics of Tuberculosis Mycobacterium, John Paul the 1st, Hospital, Krakow, Poland; ^cNantes Nanotechnological Systems, Bolesławiec, Poland; ^dFaculty of Food Technology, University of Agriculture in Krakow, Krakow, Poland

ABSTRACT

Tap water treated in air with low-temperature and low-pressure glow plasma of low frequency was tested for its either stimulation or inhibition of the growth of the selected microorganisms commonly colonizing human organism. The growth of chosen microorganisms was monitored by estimation of optical density of their colonies. The fairly linear growth against time of all microorganisms under study accelerated after 12 h from the beginning of the experiment. Colonies of *E. coli* and *S. cerevisiae* breed in the plasma treated water had an approximately 20% stimulation of the growth which was observed between 12 and 24 h. Neither stimulation nor inhibition of the growth could be noted for colonies of *Aspergillus niger*, *Candida albicans*, *Yarrowia lipolytica*, and *Enterococcus faecalis*, in whole period of observation. The plasma-treated water had no effect upon the growth of Mycobacteria. Independently of the water tested, *M. tuberculosis* started proliferating on the 14th day of the experiment, *M. intercellulare* and *M. kansasii* after 9 days, and the growth of *M. fortuitos* could be observed after 3 days.

ARTICLE HISTORY

Received 4 October 2022
Revised 19 December 2022
Accepted 20 December 2022


KEYWORDS

Aspergillus niger; *Candida albicans*; *Enterococcus faecalis*; *Escherichia coli*; *Klebsiella pneumoniae*; *Mycobacteria*; *Saccharomyces cerevisiae*; *Staphylococcus aureus*; *Yarrowia lipolytica*

Introduction

All known kinds of plasma are capable of breaking chemical bonds. They generate either/and atomic oxygen, hydrogen peroxide, and ozone. Hence, they can initiate chemical reactions.^[1–25] In contrast to them, a treatment of water with invented by Oszczyda et al.^[26, 27] low-temperature, low-pressure glow plasma of low frequency (LPGP) machine solely declusterizes the water macrostructure into smaller structural units. In this manner, the latter plasma provides a liquid with unique physical, chemical, biochemical, and functional properties.^[28] Since LPGP cannot break valence bonds, it does not initiate chemical reactions and it generates neither atomic oxygen radical nor perhydrol and ozone.

Water treated with LPGP in the air (LPGPA) better dissolves several compounds, among them also mineral salts. Simultaneously, LPGPA more readily penetrates cells of living organisms. For that sake LPGPA should be a good vector transporting such compounds to the cells. It has been proven for entomopathogenic fungi employed as biopesticides, fermentative microorganisms, and in animal breeding. Application of LPGPA appeared beneficial in agriculture and in cosmetology.^[29–37] Potentially, LPGPA could be used for preparation of various soft drinks and as media useful in human prophylaxis and therapy. However, thus far, nothing is known about the effect of LPGPA upon bacteria, fungi, and yeast-colonizing the human body. It induced these studies on the effect of LPGPA

CONTACT Jarosław Chwastowski  jaroslaw.chwastowski@pk.edu.pl  Institute of Chemistry and Inorganic Technology, Krakow University of Technology, Warszawska Str. 24, 31155 Krakow, Poland

© 2023 Jarosław Chwastowski, Katarzyna Wójcik, Henryk Kołoczek, Zdzisław Oszczyda, Karen Khachatryan and Piotr Tomasik. Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

on selected microorganisms commonly colonizing the human organism and being delivered to it. The group of selected bacteria included strains of *Escherichia coli*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, *Mycobacterium tuberculosis*, *M. intracellulare*, *M. kansasii*, and *M. fortuitum*.

Although a majority of *E. coli* strains are harmless, some strains cause harmful food poisoning.^[38, 39] The harmless strains constitute the normal microbiota of the gut, producing K₂ vitamin and benefiting their hosts, helping blood to clot, and hindering their intestines from colonization with pathogenic bacteria of a symbiotic relationship.^[40]

S. aureus, another Gram-positive bacterium, a facultative anaerobe that constitutes the microbiota of the body. It occupies the upper respiratory tracts and the skin. Turning into an opportunistic pathogen, it becomes a common cause of skin infections including abscesses and respiratory infections. Antibiotic-resistant strains of *S. aureus* have become a problem in clinical medicine.^[41]

E. faecalis, a Gram-positive, commensal bacterium, colonizes gastrointestinal tracts. It can be responsible for life-threatening infections, especially in hospitals,^[42] and reinfection of root canal-treated teeth.^[43] *K. pneumoniae*, a Gram-negative, non-motile facultative anaerobic bacterium, resides in the normal flora of the mouth, skin, and intestine. It causes destructive changes to the human lungs. It is also an important pathogen in nosocomial infections.^[42]

M. tuberculosis, an acid-fast (Ziehl-Neelsen stain), weakly Gram-positive bacterium, is an etiological agent of tuberculosis. It belongs to the family *Mycobacteriaceae* as a part of the group of mycobacteria.^[44] *Tubercle bacillus* is an aerobic (or microaerophilic) immobile, anaerobic bacterium. It does not form spores or toxins. *M. tuberculosis* is a complex of at least nine members. The cell wall of mycobacteria contains about 60% of lipids such as mycolic acids, lipoarabinomannan, and waxes, which determine its acid resistance. *M. tuberculosis* attacks lungs, kidney, spine, and brain. The disease caused by *M. tuberculosis* can be fatal.^[42]

The effect of LPGPA on a group of fungi was studied involving an *Aspergillus niger* strain producing ochratoxin A and other mycotoxins. It is one of the most abundant food-contaminating mycotoxin. The consumption of this fungus produces chronic neurotoxic, immunosuppressive, genotoxic, carcinogenic, and teratogenic effects.^[45] Its airborne spores evoke asthma in children and lung diseases.^[46]

Strains of *Candida albicans*, *Yarrowia lipolytica*, and *Saccharomyces cerevisiae* were employed to study the effect of LPGPA on yeasts. *C. albicans* is an opportunistic pathogenic yeast^[47] considered as a dimorphic fungus since it grows both as yeast and filamentous cells. It consists of several morphological phenotypes.^[48] It is common in the human gut flora and biofilms formed either on implanted medical devices or on human tissues.^[49] *C. albicans* is usually a commensal organism, but it can become pathogenic in immunocompromised individuals under a variety of conditions.^[50, 51] *S. cerevisiae*, a eukaryotic model organism, is a single-celled fungus microorganism. *S. cerevisiae* can cause invasive, life threatening infections.^[52] *Y. lipolytica* yeast species synthesize a wide group of lipases and other hydrolytic enzymes, microbial oil, citric acid, erythritol, and γ -decalactone. Although that yeast is nonpathogenic, it can induce infections in immunocompromised and critically ill patients.^[53]

In this study, the influence of LPGPA on the growth of the aforementioned microorganisms was examined. The results provided an evidence for consumers, infected with those microorganisms, whether they can safely drink LPGPA and drinks prepared on its basis. Additionally, it was evaluated for its possibility to apply LPGPA for prophylaxis and therapy.

Materials and methods

Materials

Test strains of *E. coli* ATCC 25922, *S. aureus* ATCC 25923, *E. faecalis* ATCC 19433, and *K. pneumoniae* ATCC 13883, originated from the American Type Culture Collection (Manassas,

VA, USA) and wild environmental strains, were taken from own microorganisms collection of the Cracow University of Technology. *M. tuberculosis* strains, *M. tuberculosis* H₃₇R_a ATCC 25177, *M. tuberculosis* with resistance to Rifampicin and Isoniazid, *M. intracellulare* ATCC 13950, *M. kansasii* ATCC 12478, and *M. fortuitum* ATCC 6841, were obtained from the Malopolska Central Laboratory of Tuberculosis Diagnostics, Cracow. Strain of *C. albicans* ATCC 90028 originated from the American Type Culture Collection (Manassas, VA, USA), and wild environmental strain of *C. albicans* was taken from the collection of the British National Collection of Yeast Cultures. *S. cerevisiae* DSM 1333 and *Y. lipolytica* DSM 1345 were taken from DMZS German Collection of Microorganisms and Cell Cultures GmbH in Braunschweig Germany.

Substrates

Bacteria were raised on TSA medium (trypticase soy agar) from Biomaxima, Lublin, Poland. The medium was sterilized for 15 min in an autoclave at 121°C at 2 bar. For the agar well diffusion method, TSA medium with 5 vol% ram blood was applied (Biomaxima, Lublin, Poland) according to Budzyńska et al.^[54] Fungi were raised on the Sabouraud agar medium (Biomaxima, Lublin, Poland). The medium was sterilized for 15 min in an autoclave at 121°C and pressure equal to 2 bar. For yeasts, cells were grown on the YPD medium supplemented with 2% agar (Yeast Peptone Dextrose; Merck, Germany).^[55] That medium was sterilized as above. Commercially available spring water containing 335 mg minerals/l (41 mg Ca²⁺/L, 24 mg Mg²⁺/L, 9 mg Na⁺/L, 2 mg K⁺/L, 220 mg HCO₃⁻/L, 36 mg SO₄²⁻/L, 3 mg Cl⁻/L) was used as a control.

Water treatment

Deionized water (200 mL) in 250 mL polyethylene bottles was placed in the reactor chamber^[26] and exposed to LPGP for 30 min. Plasma of 38°C was generated at 5×10^{-3} mbar, 600 V, 50 mA, and 280 GHz frequency. Treated water (LPGPA) was stored at room temperature in closed Teflon containers for no longer than one week prior to the experiments.

Breeding of microorganisms

Investigated bacteria, except *M. tuberculosis* strains, were passed onto Petri dishes with the TSA solid medium (Tryptic Soya Agar, Biomaxima, Lublin, Poland) and stored for 24 h at 44°C and 37°C. Fungi were raised for 120 h on the Petri dishes with Sabouraud solid medium (Biomaxima, Lublin, Poland) at 28°C, whereas yeasts were bread at 28°C for 72 h on the Petri dishes using solid YPD medium (Merck, Germany).

Microorganism growth

Microorganisms were sieved under a laminar flow chamber into sterilized 250 mL conical flasks containing 40 mL of liquid medium (formulated with LPGPA). The cultures were then incubated at appropriate temperature. The growth liquid medium was prepared with 1 g of peptone, 1 g of yeast extract, and 8 g of sucrose which were weighed successively into a glass screw capped bottle. Mineral salt solution was prepared by dissolving 6 g MgSO₄, 10 g K₂HPO₄, 2.4 CaCl₂, and 60 g (NH₄)₂SO₄ in 1 L distilled water. Twenty-five milliliters of that solution was added to a glass bottle containing a previously prepared liquid medium. The whole was diluted with 500 mL LPGPA and sterilized in an autoclave at 121°C for 15 min at 2 MPa.

Optical density measurements – growth evaluation

Spectrophotometric optical density (OD) measurements with a Rayleigh model UV 1800 spectrophotometer at 540 nm were carried out. The growth of microorganisms was evaluated based on the changes in the optical density in 1 mL suspensions at a given time. An increase in the solution cloudiness was measured.

Sample preparation for measuring the effect of LPGPA on *M. tuberculosis* growth

Suspensions of 0.5 McFarland density were prepared from colonies of microorganisms. Inoculum of each strain was transferred onto Lowenstein-Jensen solid media with calibrated inoculation loop. Colonies ground in a glass microbial mortar were suspended in 0.5 mL LPGPA and transferred into a sterile tube and supplemented with LPGPA to 5 mL. The density of those suspensions was measured using a densitometer and, if necessary, more LPGPA was added to achieve the desired 0.5 McFarland density. The resulting bacterial suspensions (10 μ L) after calibration with inoculation loop were inoculated onto Lowenstein-Jensen solid media. The seeded media were placed in a plastic tray and incubated at 37°C for 16 days, taking a reading every 24 h.

Statistical analysis

The experiments were conducted in triplicates. The obtained results were averaged. In all of the experiments, the observed values are given with $\pm 5\%$ standard deviations from the mean.

Results and discussion

Within the 32 h period of growth of all microorganisms some stages could be distinguished. The most frequent, occurring after a relatively fast 0–5th step growth slowed down until 24th h in order to accelerate after to the end of the experiment. Within the 32 h incubation the growth kinetics of bacterial strains of *E. coli* ATCC 25922, *S. aureus* ATCC 25923, *E. faecalis* ATCC 19433, and *K. pneumoniae* ATCC 13883 in LPGPA closely resembled that observed in the non-treated, control water (Figure 1).

In case of *E. coli* and *S. aureus*, some subtle effects of the treatment with LPGPA could be observed after 28 h LPGPA suppressed the bacteria growth to a negligible extent. The insignificant effect of LPGPA upon the growth of *E. faecalis* and *K. pneumoniae* could be observed already between 10th and 15th h. It then rose up until the 32nd h of experiment, still remaining subtle. Contact with LPGPA subtly either inhibited or stimulated the growth, respectively, depending on the test duration. LPGPA did not stimulate the growth of *M. tuberculosis* (Table 1).

In the control tests, the colonies of *M. fortuitum*, *M. intracellulare*, and *M. kansasii* and both strains of *M. tuberculosis* started rising on the 3rd, 9th, and 14th day, respectively. The contact with LPGPA did not influence that sequence. Similar responses from *A. niger*, *C. albicans*, and *Y. lipolytica* fungi showed that LPGPA was neither mycostatic nor mycotoxic (Figure 2), although between 9th and 24th h of the experiment, an approximately 10% inhibition growth could be observed for *A. niger* and *Y. lipolytica*.

C. albicans appeared insensitive to LPGPA and slight stimulation of *S. cerevisiae* was noted from the 8th day. The growth of *Y. lipolytica* and *A. niger* fungi was inhibited. Also, the growth of eukaryotic cells did not change. In order to check whether LPGPA would influence the growth of chosen microorganisms in less favorable conditions, experiments were carried out with a medium containing only mineral salts. The results were identical.

Differences in the kinetics of growth of *E. coli* and *S. cerevisiae* in the water-based minimal medium (Figure 3) revealed that LPGPA facilitated the transport of substances across the cell membranes and simultaneously, an approximately 20% stimulation of the growth could be noted in the period between




Figure 1. Comparison of prokaryotic microorganism cell growth measured as optical density (OD) in control medium (C) and plasma water-based medium (PL): A – *Escherichia coli*, B – *Staphylococcus aureus*, C – *Enterococcus faecalis*, D – *Klebsiella pneumoniae*. The tests were carried out in triplicates and averaged. Standard errors were below 5%.

Table 1. Comparison of growth of *M. tuberculosis* strains in the presence and absence of LPGPA in the culture medium^{a,b}

Microorganism	Days of growth in control 0.9% aq. NaCl solution and in LPGPA														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>M. tuberculosis</i> H37RA															X
<i>M. tuberculosis</i> MDR															X
<i>M. intracellulare</i> ATCC 13950									X						
<i>M. kansasii</i> ATCC 12478									X						
<i>M. fortitum</i> ATCC 6841			X												

^aThe tests were carried out in triplicates and averaged. Standard error was lower than 5%. ^bX – visible growth of microorganism

9th and 24th h. Inspection of Table 1 revealed that LPGPA did not influence the growth of Mycobacteria. Regardless, these microorganisms were maintained in 0.9% aq. solution of physiological salt or in LPGPA, and the observable increase in microorganisms of *M. fortuitum* took place on the third day of experiment, propagation of *M. intracellulare* and *M. kansasii* could be seen on the 9th day, and *M. tuberculosis* developed just in the 14th day of experiment (Table 1).




Figure 2. Comparison of eukaryotic cell growth measured as optical density (OD) in control medium (C) and plasma water-based medium (PL). A – *Saccharomyces cerevisiae*, B – *Candida albicans*, C – *Yarrowia lipolytica*, D – *Aspergillus niger*. The tests were carried out in triplicates and averaged. Standard errors were below 5%.




Figure 3. Comparison of cell growth in control medium (C) and plasma water-based minimal medium (PL). A – *Escherichia coli*, B – *Saccharomyces cerevisiae*. The tests were carried out in triplicates and averaged. Standard errors were below 5%.

The results showed that LPGPA did not increase the bacterial population although it facilitated the transport of substances across the cell membranes.^[29, 30] Gram positive and Gram negative bacteria showed no difference in response to the contact with LPGPA. Additional experiments were carried out with the use of a medium containing only mineral salts. Differences in kinetics of the growth of *E. coli* and *S. cerevisiae* were observed (Figure 3) supporting an assumption that LPGPA may facilitate the transport of substances across the cell membrane. Therefore, LPGPA can be safely used as potable water and for preparation of various soft drinks.

Conclusion

Water treated with low-temperature, low-pressure glow plasma of low frequency (LPGPA), regardless of the applied method of its preparation, neither inhibits nor stimulates the growth of yeast, fungi, and bacteria including *Mycobacterium* microorganisms. The results provide an evidence that the water treated with low-temperature, low-pressure glow plasma of low frequency is safe in several everyday applications, that is, for preparing various drinks and also for prophylaxis and therapy. LPGPA can be safely drunk by consumers with infectious diseases. Because LPGPA is a good vector for solutes dissolved in it, the use of drinks prepared with such kind of water can be beneficial.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

The author(s) reported that there is no funding associated with the work featured in this article.

References

- [1] Doughty, D. A.; Hartog, E. A. D.; Lawler, J. E. Current Balance at the Surface of a Cold Cathode. *Phys. Rev. Lett.* 1987, 58, 2668–2671. DOI: [10.1103/PhysRevLett.58.2668](https://doi.org/10.1103/PhysRevLett.58.2668).
- [2] Donsbach, K. W.; Cazares, R. Process for Making Highly Oxygenated Drinking Water and Drinking Water Made by the Process. U.S. Patent 5587191A, 24 December 1996.
- [3] Zelenak, Z. M.; Berzsenyi, L.; Abramo, F. Oxygen Enriched Liquids, Method and Apparatus for Making, and Applications Thereof. U.S. Patent 581422A, 29 September 1998.
- [4] Miichi, T.; Hayashi, N.; Ihara, S.; Satoh, S.; Yamade, C. Ozone Generation inside Bubbles in Water. *IEEJ Trans. Fundam. Mater.* 2001, 121, 448–452. DOI: [10.1541/ieejfms1990.121.5_448](https://doi.org/10.1541/ieejfms1990.121.5_448).
- [5] DeWald, J. J. Method and Apparatus for Adding Oxygen to Drinking Water. U.S. Patent 69361179B2, 30 August 2005.
- [6] Danone, C. G. Method for Enriching Water with Oxygen by an Electrolytic Process, Oxygen Enriched Water or Beverage and Uses Thereof. U.S. Patent 8,709,231, 19 December 2008.
- [7] Sato, K.; Yasuoka, K.; Ishii, S. Water Treatment with Pulsed Plasmas Generated inside Bubbles. *IEEJ Trans. Fundam. Mater.* 2008, 128, 401–406. DOI: [10.1541/IEEJFMS.128.401](https://doi.org/10.1541/IEEJFMS.128.401).
- [8] Verreycken, T.; Schram, D. C.; Leys, C.; Bruggeman, P. J. Spectroscopic Study of an Atmospheric Pressure Dc Glow Discharge with a Water Electrode in Atomic and Molecular Gases. *Plasma Sources Sci. Technol.* 2010, 19, 45004. DOI: [10.1088/0963-0252/19/4/045004](https://doi.org/10.1088/0963-0252/19/4/045004).
- [9] Nemcova, L.; Nikiforov, A.; Leys, C.; Krcma, F. Chemical Efficiency of $\{H\}_2\{O\}_2$ Production and Decomposition of Organic Compounds under Action of DCC under Water Discharge in Gas Bubbles. *IEEE Trans. Plasma. Sci.* 2011, 39, 865–870. DOI: [10.1109/TPS.2010.2098053](https://doi.org/10.1109/TPS.2010.2098053).
- [10] Takahashi, K.; Yagi, I.; Takaki, K.; Satta, N. Development of Pulsed Discharge inside Bubble in Water. *IEEE Trans. Plasma. Sci.* 2011, 39(11), 2654–2655. DOI: [10.1109/TPS.2011.2164095](https://doi.org/10.1109/TPS.2011.2164095).
- [11] Sommers, B.; Foster, J. Plasma Formation inside Deformed Gas Bubbles Submerged in Water. In: *Proc. Am. Phys. Soc. 65th Annual Gaseous Electronics Conference*, Austin, TX, USA, 22–26 October 2012. https://scholar.google.com/scholar_lookup?title=Plasma+formation+inside+deformed+gas+bubbles+submerged+in+water&confer

- ence=Proceedings+of+the+American+Physical+Society,+65th+Annual+Gaseous+Electronics+Conference&author=Sommers,+B.&author=Foster,+J.&publication_year=2012
- [12] Muradia, S.; Nagatsu, M. Low-voltage Pulsed Plasma Discharges inside Water Using a Bubble self-generating Parallel Plate Electrode with a Porous Ceramic. *Appl. Phys. Lett.* **2013**, *102*(14), 144105. <https://doi.org/10.1063/1.4799652>
- [13] Pranevicius, L.; Tuckute, S.; Gedvilas, K.; Gedvilas, K. Water vapor-plasma-enhanced Oxidation of Thin Titanium Films. *Acta Phys. Polon. A.* **2013**, *123*, 907–910. DOI: [10.12693/APhysPolA.123.907](https://doi.org/10.12693/APhysPolA.123.907).
- [14] Hayashi, Y.; Takada, N.; Kanda, H.; Goto, M. Generation of Pulsed Discharge Plasma in Water with Fine Bubbles. *Proc. APS Gaseous Conf.* **2015**, *LW1*, 123. https://jglobal.jst.go.jp/en/detail?GLOBAL_ID=202102254752813848
- [15] Takahashi, K.; Konno, R.; Akiyama, M.; Takaki, K.; Satta, N. Improvement of Energy Efficiency for de Colorization of Organic Dye by Discharge inside Bubble in Water, 2016. *IEEE Int. Conf. Plasma Sci. (ICOPS) 2016*, 1. DOI: [10.1109/PLASMA.2016.7533948](https://doi.org/10.1109/PLASMA.2016.7533948).
- [16] Shang, K.; Li, J.; Wang, X.; Yao, D.; Lu, N.; Jiang, N.; Wu, Y. Evaluating the Generation Efficiency of Hydrogen Peroxide in Water by Pulsed Discharge over Water Surface and Underwater Bubbling Pulsed Discharge. *Jpn. J. Appl. Phys.* **2015**, *55*(1S), 01AB02. DOI: [10.7567/JJAP.55.01AB02](https://doi.org/10.7567/JJAP.55.01AB02).
- [17] Iwabuchi, M.; Takahashi, K.; Takaki, K.; Satta, N. Influence of Sodium Carbonate on Decomposition of Formic Acid by Pulsed Discharge Plasma inside Bubble in Water. *Jpn. J. Appl. Phys.* **2016**, *55*(7S2), 07LF02. DOI: [10.7567/JJAP.55.07LF02](https://doi.org/10.7567/JJAP.55.07LF02).
- [18] Khristolubova, V. I.; Kashapov, N. F.; Shaekhov, M. F. Gas and Plasma Dynamics of RF Discharge Jet of low-pressure in a Vacuum Chamber with Flat Electrodes and inside Tube, Influence of RF Discharge on the Steel Surface Parameters. *IOP Conf. Ser. Mater. Sci. Eng.* **2016**, *134*, 012017. DOI: [10.1088/1757-899X/134/1/012017](https://doi.org/10.1088/1757-899X/134/1/012017).
- [19] Liu, J.; He, B.; Chen, Q.; Li, J.; Xiong, Q.; Yue, G.; Zhang, X.; Yang, S.; Liu, H.; Liu, Q. H. Direct Synthesis of Hydrogen Peroxide from plasma-water Interactions. *Sci. Rep.* **2016**, *6*(1), 38454. DOI: [10.1038/srep38454](https://doi.org/10.1038/srep38454).
- [20] Sharma, A.; Levko, D.; Raja, L. L.; Cha, M. S. Kinetics and Dynamics of Nanosecond Streamer Discharge in atmospheric-pressure Gas Bubble Suspended in Distilled Water under Saturated Vapour Pressure Conditions. *J. Phys. D: Appl. Phys.* **2016**, *49*, 395205. DOI: [10.1088/0022-3727/49/39/395205](https://doi.org/10.1088/0022-3727/49/39/395205).
- [21] Chen, Z.; Krasik, Y. E.; Cousens, S.; Ambujakshan, A.; Corr, C.; Dai, X. J. Generation of Underwater Discharges inside Gas Bubbles Using a 30-needles-to-plate Electrode. *J. Appl. Phys.* **2017**, *122*, 153303. DOI: [10.1063/1.4993497](https://doi.org/10.1063/1.4993497).
- [22] Takahashi, K.; Takeda, M.; Konno, R.; Takaki, K.; Satta, N. Influence of Electric Parameters on Hydroxyl Radical Production by Positive Pulsed Discharge inside of a Bubble in Water. *IEEE Trans. Plasma. Sci.* **2018**, *47*, 1105–1113. DOI: [10.1109/TPS.2018.2883767](https://doi.org/10.1109/TPS.2018.2883767).
- [23] Hagen, R. Plasma-treated Water as a Superior Electrolyte. accessed on 3 July 2017. <https://www.advancedsciencenews.com/plasma-treated-water-superior-electrolyte/>
- [24] Yui, H.; Someya, Y.; Kusama, Y.; Kanno, K.; Banno, M. Atmospheric Discharge Plasma in Aqueous Solution: Importance of the Generation of Water Vapor Bubbles for Plasma Onset and Physicochemical Evolution. *J. Appl. Phys.* **2018**, *124*, 103301. DOI: [10.1063/1.5040314](https://doi.org/10.1063/1.5040314).
- [25] Messer Americas, F. A. R. M. O. X. Drop-in Oxygenation Apparatus. New, Easy-to-deploy, Highly Efficient Solution for Oxygenation of Water. accessed on 25 June 2020. https://cdn2.hubspot.net/hubfs/189660/Messer%20US%20Website%2019/Resourses/MESS-3024_FARMOX_Dropln_datasheet.pdf
- [26] Oszczyda, Z.; Elkin, I.; Stręk, W. Equipment for Treatment of Water with Plasma Polish Patent. *Appl. PL 216025 B1*, 20.11. 2014, .
- [27] Reszke, E.; Yelkin, L.; Oszczyda, Z. Plasmating Lamp with Power Supply, Polish Patent PL 227530 B1, 2017.
- [28] Białopiotrowicz, T.; Ciesielski, W.; Domański, J.; Doscocz, M.; Fiedorowicz, M.; Grąż, K.; Khachatryan, K.; Kołoczek, H.; Kozak, A.; Oszczyda, Z., et al. Structure and Physicochemical Properties of Water Treated with low-temperature low-frequency Glow Plasma. *Curr. Phys. Chem.* **2016**, *6*, 312–320. DOI: [10.2174/18779468066661118152613](https://doi.org/10.2174/18779468066661118152613).
- [29] Tomasik, P.,. *Fundamentals of Food Nanotechnology, Essentials*. Lambert Academic Publishing: Saarbruecken, **2017**; Vol. 20017, pp 63.
- [30] Tomasik, P.,. *Fundamentals of Nanotechnology of Food and Cosmetics (In Polish)*. Sophia Scientific Editorial Board: Warsaw, **2019**; pp 54.
- [31] Szymanowicz, J.; Schwarz, T.; Murawski, M.; Małopolska, M.; Oszczyda, Z.; Tuz, R.; Nowicki, J.; Bartlewski, P. M. Storage of Bear Semen at 16–18°C in the Long Term Commercial Extender Prepared with Deionized Water or Nanowater. *Anim. Reprod.* **2019**, *164*, 1–7. DOI:[10.2478/aoas-2022-0008](https://doi.org/10.2478/aoas-2022-0008).
- [32] Ciesielska, K.; Ciesielski, W.; Kulawik, D.; Oszczyda, Z.; Tomasik, P. Cultivation of Cress Involving Water Treated under Different Atmospheres with Low-Temperature, Low- Pressure Glow Plasma of Low Frequency. *Water.* **2020**, *12*, 2152. DOI: [10.3390/w12082152](https://doi.org/10.3390/w12082152).
- [33] Ciesielska, K.; Ciesielski, W.; Girek, T.; Kołoczek, H.; Oszczyda, Z.; Tomasik, P. Reaction of *Lavandula angustifolia* Mill. to Water Treated with Low-Temperature, Low-Pressure Glow Plasma of Low Frequency. *Water.* **2020**, *12*, 3186. DOI: [10.3390/w12113168](https://doi.org/10.3390/w12113168).

- [34] Ciesielska, K.; Ciesielski, W.; Girek, T.; Oszczyda, Z.; Tomasik, P. Effect of Watering of Selected Seasoning Herbs with Water Treated with Low-temperature, Low-pressure Glow Plasma of Low Frequency. *Water*. 2020, 12, 3526. <https://doi.org/10.3390/w12123526>.
- [35] Ciesielski, W.; Gąstoł, M.; Kulawik, D.; Oszczyda, Z.; Pisulewska, E.; Tomasik, P. Specific Controlling Essential Oil Composition of Basil (*Ocimum basilicum* L.) Involving Low-temperature, Low-pressure Glow Plasma of Low Frequency. *Water*. 2020, 12, 3332. <https://doi.org/10.3390/w12123332>.
- [36] Pater, A.; Zdaniewicz, M.; Satora, P.; Khachatryan, G.; Oszczyda, Z. Application of Water Treated with Low-temperature Low-pressure Glow Plasma for Quality Improvement of Barley and Malt. *Biomolecules*. 2020, 10, 267. DOI: 10.3390/biom10020267.
- [37] Ciesielski, W.; Domagała, I.; Garcia, B.; Girek, T.; Oszczyda, Z.; Szczuka, E.; Tomasik, P. Specific Way of Controlling Composition of Cannabinoids and Essential Oil from Cannabis sativa var. Finola. *Water*. 2022, 14, 688. DOI: 10.3390/w14050688.
- [38] Tenaillon, O.; Skurnik, D.; Picard, B.; Denamur, E. The Population Genetics of Commensal *Escherichia coli*. *Nature Revs. Microbiol.* 2010, 83, 207–217. DOI:10.1038/nrmicro2298.
- [39] Center for Disease Control and Prevention. *Escherichia coli*; *Saving Lives, Protecting People*. 2020. CDC 24/7. <https://www.cdc.gov/ecoli/index.html>
- [40] Reid, G.; Howard, J.; Gan, B. S. Can Bacterial Interference Prevent Infection? *Trends Microbiol.* 2001, 9(9), 424–428. DOI: 10.1016/s0966-842x(01)02132-1.
- [41] Tong, S. Y.; Davis, J. S.; Eichenberger, E.; Holland, T. L.; Fowler, V. G. Staphylococcus aureus Infections: Epidemiology, Pathophysiology, Clinical Manifestations and Management. *Clin. Microbiol. Revs.* 2015, 28, 603–661. DOI: 10.1128/cmr.00134-14.
- [42] Ryan, K. J.; Ray, C. G. Sherris Medical Microbiology. *J. Biosci. Meds.* 2004, 3(7), 294–295.
- [43] Rocas, I.; Siqueira, J.; Santos, K. Association of *Enterococcus faecalis* with Different Forms of Periradicular Diseases. *J. Endod.* 2004, 30, 315–320. DOI: 10.1097/00004770-200405000-00004.
- [44] Virella, G.; *Microbiology and Infectious Diseases (in Polish)*. Urban & Partner: Wrocław, 2000.
- [45] Ravelo Abreu, A.; Rubio Armendáriz, C.; Gutiérrez Fernández, A. J.; Hardisson, D. L. T.; Ochratoxin, A. A in Foods for Human Consumption: Review (In Spanish). *Nutr. Hospital.* 2011, 26, 1215–1226. DOI: 10.1590/S0212-16112011000600004.
- [46] Reponen, T.; Lockey, J.; Bernstein, D. I.; Vesper, S. J.; Levin, L.; Khurana Hershey, G. K.; Zheng, S.; Ryan, P.; Grinshpun, S. A.; Villareal, M., et al. Infant Origins of Childhood Asthma Associated with Specific Moulds. *J. Allergy Clin. Immunol.* 2012, 130, 639–644.e5. DOI: 10.1016/j.jaci.2012.05.030.
- [47] Gow, N. A. R.; Yadav, B. Microbe Profile: *Candida albicans*: A shape-changing Opportunistic Pathogenic Fungus of Humans. *Microbiology.* 2017, 163, 1145–1147. DOI: 10.1099/mic.0.000499.
- [48] Basso, V.; d'Enfert, C.; Znaidi, S.; Bachellier-Bassi, S. From Genes to Networks: The Regulatory Circuitry Controlling *Candida albicans* Morphogenesis. *Curr. Topics Microbiol. Immunol.* 2019, 422, 61–99. DOI: 10.1007/82_2018_144.
- [49] Kumamoto, C. A. Candida Biofilms. *Curr. Opin.Microbiol.* 2002, 5, 608–611. DOI: 10.1016/s1369-5274(02)00371-5.
- [50] Martins, N.; Ferreira, I. C.; Barros, L.; Silva, S.; Henriques, M. Candidiasis: Predisposing Factors, Prevention, Diagnosis and Alternative Treatment. *Mycopathologia.* 2014, 177(5–6), 223–240. DOI: 10.1007/s11046-014-9749-1.
- [51] Erdogan, A.; Rao, S. S. Small Intestinal Fungal Overgrowth. *Curr. Gastroenterol. Repts.* 2015, 174, 16. DOI:10.1007/s11894-015-0436-2.
- [52] Enache-Angoulvant, A.; Hennequin, C. Invasive Saccharomyces Infection; A Comprehensive Review. *Clin. Inf. Dis.* 2005, 41, 1559–1568. DOI: 10.1086/497832.
- [53] Zienuk, B.; Fabiszewska, A. *Yarrowia lipolytica*: A Beneficial Yeast in Biotechnology as a Rare Opportunistic Fungal Pathogen: A Minireview. *World J. Microbiol. Biotechnol.* 2019, 35(1), 10. DOI: 10.1007/s11274-018-2583-8.
- [54] Budzyńska, A.; Więckowska-Szakiel, M.; Kalemba, D.; Sadowska, B.; Różalska, B. Verification of Methodical Parameters of Determination of Bactericidal Activity of Essential Oils (In Polish). *Med. Dośw. Mikrobiol.* 2009, 61, 281–287. <http://www.medmikro.org/weryfikacja-metodycznych-parametrow-oznaczania-przeciwbakteryjnej-aktywnosci-olejkow-eterycznych?lang=pl>
- [55] Atlas, R. M.; *Handbook of Microbiological Media*, 4th ed. CRC Press: Boca Raton, 2004.